

# **UKTAG River Assessment Method Macrophytes and Phytobenthos**

## **Phytobenthos – Diatoms for Assessing River and Lake Ecological Quality (River DARLEQ2)**

by

**Water Framework Directive – United Kingdom Technical Advisory Group  
(WFD-UKTAG)**



**Publisher:** **Water Framework Directive – United Kingdom Advisory Group (WFD-UKTAG)**  
c/o SEPA  
Strathallan House  
Castle Business Park  
Stirling  
FK9 4TZ  
Scotland  
[www.wfduk.org](http://www.wfduk.org)

December 2014

ISBN: 978-1-906934-58-3

**Copyright © 2014 WFD-UKTAG**

All rights reserved. WFD-UKTAG members, servants or agents accept no liability whatsoever for any loss or damage arising from the interpretation or use of the information, or reliance upon views contained herein.

**Health and safety statement**

WARNING. Working in or around water is inherently dangerous; persons using this standard should be familiar with normal laboratory and field practice. This published monitoring system does not purport to address all of the safety problems, if any, associated with its use. It is the responsibility of the user to establish appropriate health and safety practices and to ensure compliance with any national regulatory guidelines.

It is also the responsibility of the user if seeking to practise the method outlined here, to gain appropriate permissions for access to water courses and their biological sampling.

# UKTAG Guide to Phytobenthos in Rivers

## Diatoms for Assessing River and Lake Ecological Quality (DARLEQ2)

### 1 Introduction

This classification method enables the assessment of phytobenthos in rivers according to the requirements of the Water Framework Directive (WFD). Phytobenthos refers to a mostly microscopic group of organisms called algae found attached to submerged surfaces such as stones and plant stems. Assessment focuses on the diatoms, a large and diverse group of algae using a tool called “DARLEQ2” (Diatoms for **A**ssessing **R**iver and **L**ake **E**cological **Q**uality, based on a metric called the Trophic Diatom Index (TDI).

DARLEQ2 forms one part of the quality element “macrophytes and phytobenthos”. Macrophytes are assessed separately with a method called LEAFPACS2. An earlier version of DARLEQ was evaluated and revised to ensure that in combination with the revised macrophyte tool, it provides an appropriate assessment to the overall quality element of macrophytes and phytobenthos. The revised method is known as DARLEQ2.

DARLEQ2 and LEAFPACS2 results are combined to produce an overall classification for macrophytes and phytobenthos, using the worst class of either sub-element. In addition, the final combined class can be modified by taking into account the presence of bacterial tufts.

#### 1.1 Metrics

The metric used to classify phytobenthos in rivers is the TDI, and the version used in DARLEQ2 is TDI4. Diatom taxa are each assigned a score from 1 (nutrient sensitive) to 5 (nutrient tolerant) and the computed TDI4 scores range from 0 (very low nutrients) to 100 (very high nutrients). The WFD requires derivation of ecological status as an EQR (Ecological Quality Ratio). The TDI4 EQR is calculated based on observed data and predicted reference values, resulting in an overall EQR representing an ecological status class of either High, Good, Moderate, Poor or Bad. The EQR scale ranges from 0 (bad ecological status) to 1 (high ecological status).

#### 1.2 Environmental pressures to which the method is sensitive

The method is used to detect the impact of nutrient enrichment, primarily indicating response to phosphorus. However, other factors such as grazing by invertebrates and hydromorphological regimes such as flow can influence diatom abundance and composition. These may influence the overall classification result, but are not built into the classification method.

#### 1.3 Geographic application

This assessment method is appropriate for all rivers in the UK.

#### 1.4 Intercalibration

This is a process whereby all European Member States were required to compare WFD class boundary values for each biological quality element (e.g. phytobenthos, phytoplankton) to ensure similar levels are set across all countries. Once a classification method has been intercalibrated, the method must be adhered to by Member States for the purposes of WFD assessment and reporting. Intercalibration mainly focussed on the EQRs which define the

class boundaries between High and Good, and Good and Moderate. DARLEQ2 was successfully intercalibrated for rivers. The current DARLEQ2 EQR boundaries are:

High/good status	1.00
Good/moderate status	0.75
Moderate/poor status	0.50
Poor/bad status	0.25

Note that for operational purposes these boundaries are normalised to a scale of 0 – 1, as described below (section 3.1.1).

## 2 Data collection

### 2.1 Sample collection – location, frequency and sample volume

Samples of the biofilm which covers the upper surfaces of submerged cobbles or small boulders in rivers are collected by brushing or scraping with a clean toothbrush. Cobbles and small boulders are the preferred sampling substrate, but if there are no cobbles or small boulders present at the sampling site, the submerged stems of emergent macrophytes, such as *Phragmites australis*, *Sparganium erectum*, *Glyceria maxima* or *Typha* species, or leaves and stems of submerged macrophytes such as *Ranunculus* species and *Potamogeton* species are sampled instead. The sampling method is fully detailed and conforms to European Standards (CEN, 2014). Samples should be collected from mid-stream using a clean toothbrush, ensuring substrata have been submerged for at least 4 weeks prior to sampling.

#### 2.1.1 Location

Samples must be representative of conditions in the river being studied, and should be collected from habitats with similar physical conditions at all sites, especially when collecting samples on repeat visits. Samples are ideally taken from riffles, runs, glides in rivers, from sites representative of the water body as a whole, and away from obvious human impacts. Location of sites near visible discharges should be far enough downstream for the discharge to have mixed fully with the river, and one should avoid sites where direct effects of organic pollution are obvious on the invertebrate fauna. This will obviously differ between catchments, but generally a site 200-500m downstream of a discharge may be most suitable. Samples should not be taken immediately after prolonged low flows or following periods of heavy rain.

#### 2.1.2 Timing and Frequency

Two samples per year should be collected, one during spring (between March and May) and one during autumn (September to November). If this is not possible, summer sampling (June to August) is an alternative option as analysis of results has demonstrated that seasonal effects are not significant. Samples should be collected not less than two months apart.

#### 2.1.3 Sample volume

Vigorous brushing of the substrate with a clean toothbrush in a plastic tray removes the diatom film from the surface of the substrate. In situations where submerged macrophytes have been sampled, cut random lengths are put into clean sampling bottles or bags and

vigorously agitated to dislodge the attached diatoms. This process results in a nominal volume of sample collected that is easily transferred to a plastic container (volume 50-100mls). There is not a precise volume of sample required, but larger volumes should be avoided.

## **2.2 Sample analysis**

If samples cannot be analysed soon after collection they should be preserved as soon as possible with Lugol's iodine (adding approximately 5-10% by volume). Samples are then digested to remove all internal contents of the diatom cell (the frustule), leaving clean 'valves' (digestion of the frustule generally results in separation of two valves). Permanent slides are prepared using Naphrax as a diatom mountant, and at least 300 undamaged valves of non-planktic taxa should be identified and counted using a high power microscope (x1000 magnification). The presence and number of valves is recorded of each diatom taxa present. The analytical method is fully detailed and conforms to European Standards (CEN, 2014).

## **2.3 Other data requirements**

Alkalinity data are required as a predictor of reference (expected) conditions for the DARLEQ2 EQR. Data should be obtained from analysis of samples from the water body taken at monthly intervals over a period of at least one year, reported as mg/L CaCO<sub>3</sub>. Alkalinity is the observed annual mean alkalinity of the site unless the observed annual mean alkalinity is <5 mg L<sup>-1</sup> CaCO<sub>3</sub>, in which case the value is set at "5"; or ≥250 mg L<sup>-1</sup> CaCO<sub>3</sub>, in which case the value is set at "250". It is recommended that the same alkalinity value is used as a predictor variable for both macrophyte and phytobenthos classifications.

Analysis carried out for the UKTAG (UKTAG, 2013) has confirmed that a DARLEQ2 based assessment alone would generally give a reliable classification if alkalinity is < 75 mg L<sup>-1</sup> CaCO<sub>3</sub> (because the diatom-based status is consistently lower than the macrophyte-based status over this range of alkalinity and will thus define the overall classification), whilst a LEAFPACS2 based assessment alone is adequate at >200 mg L<sup>-1</sup> CaCO<sub>3</sub> (because the macrophyte-based status is then consistently lower than the diatom-based status). In the middle range, both components are necessary (because neither one is consistently lower than the other) although, on average, an assessment based on DARLEQ2 alone will be more reliable guide than LEAFPACS2 alone at alkalinities up to ~120 mg L<sup>-1</sup> CaCO<sub>3</sub>, whereas LEAFPACS2 alone will be a more reliable guide at alkalinities above this.

### **2.3.1 Bacterial tufts**

The normative definition in the WFD (Annex V) refers to the "displacement" of macrophytes and phytobenthos by bacterial tufts and coats at moderate status, implying a need to recognise a state where the organic loading is so high that heterotrophic organisms can out-compete phototrophic organisms. Bacterial tufts and coats are generally referred to as "sewage fungus", which include a mixture of heterotrophic bacteria, fungi and protozoans.

Presence of sewage fungus should be recorded in the field on all occasions a diatom sample is collected, recording cover: occasional (<30% of surface area); widespread (30-60% of surface area) or extensive (>61% of surface area), and density: trace (but only just detectable); thin (obvious presence but substrate not obscured); thick (thick enough to fully obscure substrate) or massive (occupies a significant proportion of the water column). Sewage fungus should only have the potential to downgrade from Moderate or worse class.

Currently UK agencies implement this where appropriate. A weight of evidence approach is adopted, where sewage fungus coverage and density (as an average over the assessment period) should over-ride the results of LEAFPACS and DARLEQ assessments to determine final status. This occurs in situations where LEAFPACS and DARLEQ assessments suggest status is Moderate or lower, and where the sewage fungus assessment suggests a lower class than the combined LEAFPACS and DARLEQ assessment.

#### **2.4 *Minimum data requirements***

Confidence of classification will depend on the number of samples taken, and the proximity of the resulting EQR to a class boundary. Ideally a total of 6 samples (over 3 years) are recommended. In practice fewer are used to produce a classification, although with a reduced confidence of class.

#### **2.5 *Typology***

Typology defines waterbodies by factors which have a strong influence on their ecology; for calculation of river phytobenthos EQR, sites require input of annual mean alkalinity as detailed above.

### 3 Procedures for calculating metric EQR

The phytobenthos EQR is a ratio of observed to expected values. *Observed values* are taken from samples collected in the field and *expected values* (reference values) are predicted from alkalinity.

#### 3.1 Calculating EQR for DARLEQ

##### 3.1.1 Status class boundaries

To enable a confidence of class to be calculated for all 5 status classes it is necessary to make an arithmetic adjustment to the intercalibrated EQR boundary values, and the calculated EQRs, so that EQR values are constrained to a scale of 0 – 1. This is achieved by multiplying the intercalibrated boundary values, and the calculated EQR, by 0.8. Thus the operational boundary values are as shown below:

	Intercalibrated boundary	Normalised operational boundary
High/Good	1.00	0.80
Good/Moderate	0.75	0.60
Moderate/Poor	0.50	0.40
Poor/Bad	0.25	0.20

This procedure does not alter the classification outcomes, it simply changes the scale on which the class boundary values are calculated and assessed.

##### 3.1.2 Manual calculation of EQR and status class

Calculation of phytobenthos EQR can be carried out manually as detailed below.

All diatom taxa are assigned a score from 1 (nutrient sensitive) to 5 (nutrient tolerant). The list of diatom taxa and their nutrient sensitivity scores is detailed in Appendix A.

The observed TDI for each sample is calculated using equations 1 and 2:

$$\begin{array}{l} \text{Observed value of} \\ \text{river trophic diatom} \\ \text{index} \end{array} = (W \times 25) - 25 \quad \text{Equation 1}$$

where:

"W" is given by the equation:

$$W = \frac{\sum_{j=1}^n a_j \times s_j}{\sum_{j=1}^n a_j} \quad \text{Equation 2}$$

where:

"a<sub>j</sub>" is the number of valves of taxon j, and

"s<sub>j</sub>" is the nutrient sensitivity score in Appendix A corresponding to the taxon represented by j. The expected (reference) value of the TDI (eTDI) is predicted from a regression equation derived from a subset of reference sites characterized by very low levels of human pressure. The model uses alkalinity, an environmental variable that reflects background geology and fertility.

The value for the expected TDI (eTDI) at reference conditions applicable to the river should be calculated using the following equation:

$$eTDI = 9.933 * \text{Exp}(\text{Log}_{10}(\text{alkalinity}) * 0.81) \quad \text{Equation 3}$$

where:

"alkalinity" is the observed annual mean alkalinity of the river in mg/l CaCO<sub>3</sub> unless the observed annual mean alkalinity is < 5 mg L<sup>-1</sup> CaCO<sub>3</sub>, in which case the value is set at "5"; or ≥250 mg L<sup>-1</sup> CaCO<sub>3</sub>, in which case the value is set at "250".

The EQR is calculated using the following equation:

$$EQR = \frac{\text{observed value of river trophic diatom index}}{(100 - \text{river trophic diatom index})} \div \frac{\text{reference value for river trophic diatom index}}{(100 - \text{river trophic diatom index})}$$

Equation 4

The final EQR value for each sample is then normalized so that it conforms to a scale from 0-1. To normalize the EQR, multiply the EQR value by 0.8. This normalization step allows the confidence of class to be calculated correctly.

Where the calculated EQR is > 1.0, its value should be set to "1".

### 3.1.3 Worked example

The following details a manual worked example from an analysis of a real sample collected from a site in the English midlands:

Taxon name	Abundance (a = number of valves)	Nutrient sensitivity score (s)	a x s
Achnanthydium minutissimum type	12	2	24
Amphora montana	1	5	5
Amphora pediculus	1	5	5
Cocconeis pediculus	5	4	20
Cocconeis placentula	61	3.1	189.1
Cyclotella atomus	1	0	0
Cyclotella meneghiniana	3	0	0



Taxon name	Abundance (a = number of valves)	Nutrient sensitivity score (s)	a x s
Diatoma vulgare	1	5	5
Encyonema minutum (E. "ventricosum")	1	2.6	2.6
Encyonema silesiacum (E. "ventricosum")	14	2.6	36.4
Fistulifera / Mayamaea	1	3.9	3.9
Fragilaria vaucheriae	1	2	2
Gomphonema "intricatum" type	3	3.6	10.8
Gomphonema clavatum	1	2	2
Gomphonema parvulum	3	3	9
Melosira varians	3	4	12
Navicula capitatoradiata	54	4	216
Navicula cryptotenella	107	4	428
Navicula gregaria	2	4	8
Navicula lanceolata	1	4	4
Navicula menisculus	1	4	4
Navicula minima	1	3	3
Navicula reichardtiana	1	4	4
Navicula tripunctata	2	5	10
Nitzschia fonticola	2	4	8
Nitzschia gracilis	1	3	3
Nitzschia palea	7	4	28
Nitzschia paleacea	3	3	9
Nitzschia sp.	1	3	3
Planothidium frequentissimum	2	3	6
Reimeria sinuata ("Reimeria sp.")	6	3	18
Reimeria uniseriata ("Reimeria sp.")	4	3	12
Rhoicosphenia abbreviata	7	4	28
Unidentified pennate diatom	1	3	3
$\Sigma a =$	315	$\Sigma as =$	1121.8

The observed value for the river trophic diatom index for this sample is calculated as follows:

sum as for all taxa in sample = 1121.8

sum a for all taxa in sample = 315

$$\text{Calculate } W = \text{sum as} / \text{sum a} = 3.56$$

Calculate the observed value of TDI using equations 1 and 2:

$$(W \times 25) - 25 = (3.74 \times 25) - 25 = 64.0$$

The mean total alkalinity for this sample was 88.5 mg/l CaCO<sub>3</sub>.

The reference value for river TDI4 for the sample is:

$$9.933 * \text{Exp}(\text{Log}_{10}(88.5)*0.81) = 48.1$$

The EQR for the sample is:

$$(100 - 64.0) / (100 - 48.1) = 36 / 51.9 = 0.69$$

Normalise the EQR by multiplying by 0.8 to give a final EQR of

$$0.69 \times 0.8 = 0.552$$

An EQR of 0.552 gives a face value class of **Moderate** status (using the normalised boundary values).

An average of EQRs obtained over a year provides an annual EQR.

#### 3.1.4 Automated calculation of EQR and status class

An MS Excel spreadsheet has been produced with a series of worksheets for data input and calculation of all components of the classification, including Confidence of Class.

Instructions for use are provided within the spreadsheet calculator. The most recent version of the spreadsheet calculator can be found on the UKTAG website. Automated data entry and calculation of diatom EQR using the DARLEQ2 calculator spreadsheet is recommended for large data sets.

*NB As revisions are likely to be made and the calculator updated over time, it is important to check that the most recent version is being used.*

#### **4 Procedures for calculating statistical confidence in metric**

The assessment of statistical error associated with each EQR entry is calculated within DARLEQ2 and is expressed as a “confidence of class”, i.e. the statistical confidence we have of the metric falling into each of the five classes, from High to Bad. This also makes it possible to determine the statistical confidence of the river classifying as “worse than Good status”.

#### **5 References**

CEN (2014). *Water quality - Guidance standard for the routine sampling and preparation of benthic diatoms from rivers and lakes*. EN 13946: 2014. Comité European de Normalisation, Geneva.

CEN (2014). *Water quality – Guidance for the identification and enumeration of benthic diatom samples from rivers and lakes*. EN 14407: 2014. Comité European de Normalisation, Geneva.

Environment Agency (2012). *A streamlined taxonomy for the Trophic Diatom Index*. Environment Agency Report SC070034/TR1.

UKTAG (2013) *Final recommendations on new and updated biological standards*. September 2013. UKTAG 2013.

[http://www.wfduk.org/sites/default/files/Media/UKTAG%20Final%20recommendations%20on%20biological%20stds\\_20131030.PDF](http://www.wfduk.org/sites/default/files/Media/UKTAG%20Final%20recommendations%20on%20biological%20stds_20131030.PDF)

## Appendix A

The following table lists the UK streamlined diatom taxa where recorded maximum relative abundance was  $\geq 2\%$  (EA, 2012) for use in TDI4.

Taxa name	Sensitivity score
<i>Achnanthes clevei</i>	5
<i>Achnanthes exigua</i>	3
<i>Achnanthes oblongella</i>	1
<i>Achnanthes</i> sp.	3
<i>Achnantheidium eutrophilum</i>	2
<i>Achnantheidium microcephalum</i> .	1
<i>Achnantheidium minutissimum</i>	2
<i>Achnantheidium pyrenaicum</i>	2
<i>Achnantheidium</i> sp..	2
<i>Achnantheidium subatomus</i>	2
<i>Adlafia minuscula</i>	3
<i>Adlafia minuscula</i> var. <i>muralis</i>	5
<i>Amphipleura pellucida</i>	3
<i>Amphora libyca</i> .	4
<i>Amphora montana</i>	5
<i>Amphora ovalis</i>	5
<i>Amphora pediculus</i> agg.	5
<i>Amphora</i> sp..	4
<i>Amphora veneta</i> .	5
<i>Bacillaria paradoxa</i>	5
<i>Brachysira brebissonii</i>	1
<i>Brachysira vitrea</i>	1
<i>Caloneis bacillum</i>	4
<i>Caloneis hyalina</i>	5
<i>Caloneis silicula</i>	4
<i>Caloneis</i> sp.	3
<i>Cavinula cocconeiformis</i>	2
<i>Cavinula variostrata</i>	3
<i>Cocconeis diminuta</i>	5
<i>Cocconeis pediculus</i>	4
<i>Cocconeis placentula</i>	3.1
<i>Cocconeis scutellum</i>	3
<i>Cocconeis</i> sp.	3
<i>Craticula accomoda</i>	4
<i>Craticula halophila</i>	4
<i>Craticula molestiformis</i>	5
<i>Ctenophora pulchella</i>	2
<i>Cymatopleura solea</i>	5
<i>Cymbella affinis</i>	2
<i>Cymbella cistula</i>	2
<i>Cymbella helvetica</i>	2

Taxa name	Sensitivity score
<i>Cymbella lanceolata</i>	4
<i>Cymbella</i> sp.	3
<i>Cymbellonitzschia diluviana</i>	4
<i>Delicata delicatula</i>	1
<i>Denticula tenuis</i>	2
<i>Diadesmis contenta</i>	5
<i>Diadesmis contenta</i> fo. <i>biceps</i>	3
<i>Diadesmis gallica</i>	3
<i>Diatoma ehrenbergii</i>	1
<i>Diatoma mesodon</i>	1
<i>Diatoma moniliformis</i>	1
<i>Diatoma problematica</i>	2
<i>Diatoma</i> sp.	2
<i>Diatoma tenue</i>	1
<i>Diatoma vulgare</i>	5
<i>Didymosphenia geminata</i>	1
<i>Diploneis elliptica</i>	5
<i>Diploneis marginestriata</i>	5
<i>Diploneis oblongella</i>	5
<i>Ellerbeckia arenaria</i>	4
<i>Encyonema 'ventricosum' agg.</i>	2.6
<i>Encyonema caespitosum</i>	4
<i>Encyonema gracile</i>	1
<i>Encyonema hebridicum</i>	1
<i>Encyonema prostratum</i>	5
<i>Encyonema</i> sp.	4
<i>Encyonopsis cesatii</i>	2
<i>Encyonopsis falaisensis</i>	1
<i>Encyonopsis microcephala</i>	2
<i>Eolimna minima</i>	3
<i>Eolimna subminuscula</i>	4
<i>Eolimna submuralis</i>	5
<i>Epithemia adnata</i>	5
<i>Epithemia sorex</i>	3
<i>Epithemia</i> sp.	2
<i>Epithemia turgida</i>	1
<i>Eucocconeis flexella</i>	1
<i>Eucocconeis laevis</i>	1
<i>Eunotia</i> sp.	1.4
<i>Fallacia helensis</i>	5
<i>Fallacia indifferens</i>	2

Taxa name	Sensitivity score
<i>Fallacia lenzii</i>	5
<i>Fallacia pygmaea</i>	5
<i>Fallacia subhamulata</i>	5
<i>Fistulifera/Mayamaea</i> spp.	3.9
<i>Fragilaria bidens</i>	3
<i>Fragilaria capucina</i>	1.2
<i>Fragilaria fasciculata</i>	4
<i>Fragilaria nitzschioides</i>	3
<i>Fragilaria</i> sp.	2
<i>Fragilaria vaucheriae</i>	2
<i>Fragilariforma</i> sp.	2.5
<i>Frustulia krammeri</i>	1
<i>Frustulia</i> sp.	5
<i>Frustulia vulgaris</i>	2
<i>Geissleria acceptata</i>	4
<i>Geissleria ignota</i>	4
<i>Geissleria schoenfeldii</i>	3
<i>Gomphonema 'intricatum'</i> type	3.6
<i>Gomphonema acuminatum</i>	2
<i>Gomphonema angustatum</i>	3
<i>Gomphonema anoenum</i>	4
<i>Gomphonema clavatum</i>	2
<i>Gomphonema clevei</i>	1
<i>Gomphonema constrictum</i> var. <i>capitatum</i>	2
<i>Gomphonema gracile</i>	1
<i>Gomphonema insigne</i>	5
<i>Gomphonema olivaceoides</i>	2
<i>Gomphonema olivaceum</i>	3
<i>Gomphonema parvulum</i>	3
<i>Gomphonema pseudoaugur</i>	5
<i>Gomphonema</i> sp.	3
<i>Gomphonema tergestinum</i>	4
<i>Gomphonema truncatum</i>	3
<i>Gomphonema ventricosum</i>	1
<i>Gomphosphenia grovei</i>	4
<i>Gyrosigma acuminatum</i>	5
<i>Gyrosigma attenuatum</i>	5
<i>Gyrosigma nodiferum</i>	4
<i>Gyrosigma scalproides</i>	5
<i>Hannaea arcus</i>	1
<i>Hantzschia abundans</i>	5
<i>Hantzschia amphioxys</i>	4
<i>Karayevia laterostrata</i>	3
<i>Kolbesia ploenensis</i>	5
<i>Lemnicola hungarica</i>	5

Taxa name	Sensitivity score
<i>Luticola goeppertiana</i>	5
<i>Luticola mutica</i>	3
<i>Luticola</i> sp.	4
<i>Luticola ventricosa</i>	3
<i>Melosira varians</i>	4
<i>Meridion circulare</i>	2
<i>Meridion circulare</i> var. <i>constrictum</i>	1
<i>Navicula</i> [small species]	4
<i>Navicula angusta</i>	1
<i>Navicula capitata</i>	4
<i>Navicula capitatoradiata</i>	4
<i>Navicula cari</i>	4
<i>Navicula carteri</i>	3
<i>Navicula cincta</i>	4
<i>Navicula claytonii</i>	3
<i>Navicula cryptocephala</i>	3
<i>Navicula cryptotenella</i>	4
<i>Navicula digitoradiata</i>	4
<i>Navicula gregaria</i>	4
<i>Navicula hungarica</i>	5
<i>Navicula integra</i>	4
<i>Navicula lanceolata</i>	4
<i>Navicula menisculus</i>	4
<i>Navicula modica</i>	5
<i>Navicula oblonga</i>	4
<i>Navicula radiosa</i>	3
<i>Navicula recens</i>	5
<i>Navicula reichardtiana</i>	4
<i>Navicula reinhardtii</i>	4
<i>Navicula rhynchocephala</i>	2
<i>Navicula schroeterii</i>	5
<i>Navicula slesvicensis</i>	3
<i>Navicula</i> sp.	4
<i>Navicula subrhynchocephala</i>	4
<i>Navicula subrotundata</i>	5
<i>Navicula tenelloides</i>	4
<i>Navicula tripunctata</i>	5
<i>Navicula trivialis</i>	4
<i>Navicula veneta</i>	4
<i>Navicula viridula</i>	4
<i>Nitzschia acicularioides</i>	5
<i>Nitzschia acicularis</i>	3
<i>Nitzschia amphibia</i>	5
<i>Nitzschia archibaldii</i>	2
<i>Nitzschia brevissima</i>	2

Taxa name	Sensitivity score
<i>Nitzschia capitellata</i>	4
<i>Nitzschia clausii</i>	4
<i>Nitzschia communis</i>	5
<i>Nitzschia disputata</i>	2
<i>Nitzschia dissipata</i>	3
<i>Nitzschia dissipata</i> subsp. <i>media</i>	3
<i>Nitzschia filiformis</i>	5
<i>Nitzschia flexa</i>	3
<i>Nitzschia fonticola</i>	4
<i>Nitzschia fossilis</i>	5
<i>Nitzschia frustulum</i>	3
<i>Nitzschia gracilis</i>	3
<i>Nitzschia hantzschiana</i>	2
<i>Nitzschia heufleriana</i>	4
<i>Nitzschia inconspicua</i>	4
<i>Nitzschia lacuum</i>	2
<i>Nitzschia liebetruthii</i>	1
<i>Nitzschia linearis</i>	4
<i>Nitzschia littoralis</i>	4
<i>Nitzschia microcephala</i>	3
<i>Nitzschia palea</i>	4
<i>Nitzschia paleacea</i>	3
<i>Nitzschia paleaeformis</i>	3
<i>Nitzschia perminuta</i>	3
<i>Nitzschia perspicua</i>	3
<i>Nitzschia pusilla</i>	4
<i>Nitzschia recta</i>	4
<i>Nitzschia sigma</i>	4
<i>Nitzschia sigmoidea</i>	3
<i>Nitzschia sociabilis</i>	4
<i>Nitzschia</i> sp.	3
<i>Nitzschia subacicularis</i>	4
<i>Nitzschia sublinearis</i>	4
<i>Nitzschia supralitorea</i>	5
<i>Nitzschia tubicola</i>	4
<i>Nitzschia vermicularis</i>	4
<i>Parlibellus protracta</i>	4
<i>Peronia fibula</i>	1
<i>Pinnularia</i> sp.	2.2
<i>Placoneis clementis</i>	4
<i>Placoneis elginensis</i>	5
<i>Planothidium bioretti</i>	2
<i>Planothidium delicatulum</i>	5
<i>Planothidium dubium</i>	3
<i>Planothidium ellipticum</i>	3

Taxa name	Sensitivity score
<i>Planothidium frequentissimum</i>	3
<i>Planothidium granum</i>	5
<i>Planothidium lanceolatum</i>	4
<i>Planothidium rostratum</i>	5
<i>Planothidium</i> sp.	4
<i>Platessa conspicua</i>	5
<i>Psammothidium chlidanos</i>	2
<i>Psammothidium grishunun</i> fo. <i>daonensis</i>	2
<i>Psammothidium helveticum</i>	2
<i>Psammothidium lauenburgianum</i>	5
<i>Psammothidium levanderi</i>	3
<i>Psammothidium marginulatum</i>	2
<i>Psammothidium scoticum</i>	2
<i>Psammothidium</i> sp.	2
<i>Psammothidium subatomoides</i>	2
<i>Pseudostaurosira/Staurosira</i> agg.	3.7
<i>Reimeria</i> sp.	3
<i>Rhoicosphenia abbreviata</i>	4
<i>Rossithidium</i> sp.	1
<i>Sellaphora joubaudii</i>	4
<i>Sellaphora pupula</i>	4
<i>Sellaphora seminulum</i>	4
<i>Simonsenia delognei</i>	5
<i>Stauroneis anceps</i>	2
<i>Stauroneis kriegeri</i>	2
<i>Stauroneis</i> sp.	3
<i>Stauroneis thermicola</i>	3
<i>Staurosirella leptostauron</i>	4
<i>Staurosirella pinnata</i>	4
<i>Surirella angusta</i>	3
<i>Surirella brebissonii</i>	3
<i>Surirella crumena</i>	3
<i>Surirella islandica</i>	3
<i>Surirella linearis</i>	1
<i>Surirella minuta</i>	4
<i>Surirella ovalis</i>	5
<i>Surirella roba</i>	2
<i>Surirella</i> sp.	2
<i>Surirella terricola</i>	4
<i>Synedella parasticia</i>	5
<i>Synedra acus</i>	3
<i>Synedra famelica</i>	2
<i>Synedra tenera</i>	1
<i>Synedra ulna</i>	2
<i>Tabellaria</i> sp.	1

<i>Taxa name</i>	Sensitivity score
<i>Tryblionella acuminata</i>	5
<i>Tryblionella apiculata</i>	5
<i>Tryblionella debilis</i>	4

<i>Taxa name</i>	Sensitivity score
<i>Tryblionella hungarica</i>	4
<i>Tryblionella sp.</i>	4